Quantifying methane emissions from the largest oil-producing basin in the United States from space

Yuzhong Zhang (Zhangyuzhong@westlake.edu.cn)

With

Ritesh Gautam, Sudhanshu Pandey, Mark Omara, Joannes D. Maasakkers, Pankaj Sadavarte, David Lyon, Hannah Nesser, Melissa P. Sulprizio,Daniel J. Varon, Ruixiong Zhang, Sander Houweling, Daniel Zavala-Araiza, Ramon A. Alvarez, Alba Lorente, Steven P. Hamburg, Ilse Aben, Daniel J. Jacob

Methane: a potent greenhouse gas

Methane concentration in last 35 years



Methane concentration in last 300 years



Radiative forcing



IPCC AR5

Reduce methane emissions

Necessary supplement to fast CO_2 reduction to achieve climate goals



IPCC, 2018

Economical, technological, & legal feasibility

Global Methane Initiative

OGCI OIL AND GAS CLIMATE INITIATIVE

METHANE GUIDING PRINCIPLES

Challenging demands for monitoring

Global coverage
Emissions from every corner of the world counts

Monitor emissions
Large numbers, varied kinds, distributed over globe, highly variable

Integrate information across scales
From global budget, regional hotspots to fixing malfunctioning facility

Global/National



5-10 km resolution
~1000 km swath

Regional



~100-1000 m resolution
~100 km swath

Facility Scale



GHGSat 8/7/18,
Permian

~10 m resolution
~10 km swath

Rapidly expanding satellite capability





Rapidly expanding satellite capability





Use satellite observations to quantify methane emissions





$$J(\boldsymbol{x}) = (\boldsymbol{x} - \boldsymbol{x}_a)^T \mathbf{S}_a^{-1} (\boldsymbol{x} - \boldsymbol{x}_a) + \gamma (\boldsymbol{y} - \mathbf{K}\boldsymbol{x})^T \mathbf{S}_0^{-1} (\boldsymbol{y} - \mathbf{K}\boldsymbol{x})$$

Prior

Prior uncertainty

Satellite

Instrument & model uncertainty

Forward model

Methane emissions from the Permian Basin

Tropospheric Monitoring Instrument (TROPOMI)



Satellite: Sentinel-5 Precursor; Swath width: 2600 km;

Overpass: ~13:30 LT; Resolution: 7×7 km^2; Retrieval: “full physics” (*Hu et al., 2016*)



Zhang et al., Science Advances, 2020

Permian Basin

Permian Basin: the largest oil producing basin in the U.S.
a lack of “top-down” constraint for its methane emissions

REPORT

Assessment of methane emissions from the U.S. oil and gas supply chain

Ramón A. Alvarez[1,*], Daniel Zavala-Araiza[1], David R. Lyon[1], David T. Allen[2], Zachary R. Barkley[3], Adam R. Brandt[4], Kenneth J. Dav…
+ See all authors and affiliations

Science 13 Jul 2018:
Vol. 361, Issue 6398, pp. 186-188
DOI: 10.1126/science.aar7204



Alvarez et al., Science, 2018

Permian Basin



Oil & gas production



Methane emissions from the Permian Basin

Methane concentration

TROPOMI

5/2018-3/2019

Solve inversion

Methane flux



Spatial distribution



Temporal changes



Spatial distributions of methane emissions

TROPOMI inversion



Oil production



Natural gas production





Estimate methane emissions with multiple approaches

Based on TROPOMI data

Atmospheric inverse modeling

0.25x0.3125 GEOS-Chem nested

Yuzhong Zhang (Westlake)



Mass balance method

Sudhanshu Pandey (SRON)

→ 3.2±2.0 Tg a^{-1}

Schneising et al., ACP, 2020

Based on surface measurements

Site-level measurement extrapolation

71 site-level measurements

Mark Omara (EDF)

Robertson et al., EST, 2020



Based on bottom-up information

Bottom-up emission inventory

Extrapolation of EPA gridded inventory to 2018 DI info for O&G

Bram Maasakkers (SRON)



More than 2 times the bottom-up estimate



>2 times the bottom-up estimate
4x higher than Eagle Ford -- the largest flux reported in literature

High leakage rate

Leakage rate
with respect to gas production



Leakage rate vs gas production



High gas production & high leakage rate indicates low efficiency in gas utilization

Intensive gas flaring over the Permian

TROPOMI inversion



2.9 Tg a^{-1}

VIIRS Infrared radiation



TROPOMI NO_2



Burning of 5.2 – 8.7 10^9 m^3 a^{-1} natural gas
→ ~ 3 Tg a^{-1}

Global emission and trend

GOSAT （2010-2018）



Averaging kernel sensitivities

Surface



Satellite





Zhang et al., ACPD, 2020; Lu et al., ACPD, 2020; Maasakkers et al., ACP, 2019

Global emission and trend

GOSAT （2010-2018）





Zhang et al., ACPD, 2020



Maasakkers et al., ACPD, 2020

Summary

Yuzhong Zhang
(zhangyuzhong@westlake.edu.cn)

- Demonstrate the capability of satellite observations to constrain regional methane emissions, through a case study at the Permian Basin
- Inverse analysis infers an annual emissions of 2.7 Tg a^{-1} from the Permian Basin, more than 2 times the bottom-up estimate
- Low rate in gas capture/utilization contributes to high leakage rate
- Globally, satellite observations supplement surface observations for understanding global methane budget and its changes.